

INFLATION

G.LAZARIDES

*Physics Division , School of Technology, Aristotle University of Thessaloniki,
Thessaloniki GR 540 06, Greece.*

The shortcomings of the Standard Big Bang Cosmological Model as well as their resolution in the context of inflationary cosmology are discussed. The inflationary scenario and the subsequent oscillation and decay of the inflaton field are then studied in some detail. The density perturbations produced during inflation and their evolution during the matter dominated era are presented. The temperature fluctuations of the cosmic background radiation are summarized. The non-supersymmetric as well as the supersymmetric hybrid inflationary model is introduced and the ‘reheating’ of the universe is analyzed in the context of the latter and a left-right symmetric gauge group. The scenario of baryogenesis via a primordial leptogenesis is considered in some detail. It is, finally, pointed out that, in the context of a supersymmetric model based on a left-right symmetric gauge group, hybrid inflation, baryogenesis via primordial leptogenesis and neutrino oscillations are linked. This scheme, supplemented by a familiar ansatz for the neutrino Dirac masses and mixing of the two heaviest families and with the MSW resolution of the solar neutrino puzzle, implies that $1 \text{ eV} \lesssim m_{\nu\tau} \lesssim 9 \text{ eV}$. The mixing angle $\theta_{\mu\tau}$ is predicted to lie in a narrow range which will be partially tested by the Chorus/Nomad experiment.

1 Shortcomings of the Big Bang Model

The Standard Big Bang (SBB) Cosmological Model¹ has been very successful in explaining, among other things, the Hubble expansion of the universe, the existence of the Cosmic Background Radiation (CBR) and the abundances of the light elements which were formed during primordial nucleosynthesis. Despite its great successes, this model had a number of long-standing shortcomings which we will now summarize:

1.1 Horizon Problem

The CBR, which we receive now, was emitted at the time of ‘decoupling’ of matter and radiation (which essentially coincides with the time of recombination of atoms) when the cosmic temperature was $T_d \approx 3,000 \text{ K}$. The decoupling time, t_d , can be calculated from

$$\frac{T_0}{T_d} = \frac{2.73 \text{ K}}{3,000 \text{ K}} = \frac{a(t_d)}{a(t_0)} = \left(\frac{t_d}{t_0} \right)^{2/3}, \quad (1)$$

where t_0 , T_0 are the present cosmic time and temperature of CBR and $a(t)$ is the dimensionless scale factor of the universe at cosmic time t normalized

so that $a(t_0) = 1$. It turns out that $t_d \approx 200,000 h^{-1}$ years, where h is the present value of the Hubble parameter in units of $100 \text{ km sec}^{-1} \text{ Mpc}^{-1}$.

The distance over which the photons of the CBR have travelled since their emission is

$$a(t_0) \int_{t_d}^{t_0} \frac{dt'}{a(t')} = 3t_0 \left[1 - \left(\frac{t_d}{t_0} \right)^{2/3} \right] \approx 3t_0 \approx 6,000 h^{-1} \text{ Mpc} , \quad (2)$$

which essentially coincides with the present particle horizon size. A sphere around us with radius equal to this distance is called the ‘last scattering surface’ since the CBR observed now has been emitted from it. The particle horizon size at t_d was $2H^{-1}(t_d) = 3t_d \approx 0.168 h^{-1} \text{ Mpc}$ ($H(t)$ being the Hubble parameter at cosmic time t) and expanded till now to become $0.168 h^{-1}(a(t_0)/a(t_d)) \text{ Mpc} \approx 184 h^{-1} \text{ Mpc}$. The angle subtended by this ‘decoupling’ horizon at present is $\theta_d \approx 184/6,000 \approx 0.03 \text{ rads} \approx 2^\circ$. Thus, the sky splits into $4\pi/(0.03)^2 \approx 14,000$ patches that never communicated causally before sending light to us. The question then arises how come the temperature of the black body radiation from all these patches is so accurately tuned as the measurements of the Cosmic Background Explorer² (COBE) require ($\delta T/T \approx 6.6 \times 10^{-6}$).

1.2 Flatness Problem

The present energy density, ρ , of the universe has been observed to lie in the relatively narrow range $0.1\rho_c \lesssim \rho \lesssim 2\rho_c$, where ρ_c is the critical energy density corresponding to a flat universe. The lower bound has been derived from estimates of galactic masses using the virial theorem whereas the upper bound from the volume expansion rate implied by the behavior of galactic number density at large distances. The Friedmann equation

$$H^2 = \frac{8\pi G}{3}\rho - \frac{k}{a^2} = \frac{8\pi G}{3}\rho_c , \quad (3)$$

where $H = \dot{a}(t)/a(t)$ (overdots denote derivation with respect to cosmic time) is the Hubble parameter, G the Newton’s constant and k negative zero or positive for an open, flat or closed universe respectively, implies that $(\rho - \rho_c)/\rho_c = 3(8\pi G\rho_c)^{-1}(k/a^2)$ is proportional to a , for matter dominated universe. Consequently, in the early universe, $|(\rho - \rho_c)/\rho_c| \ll 1$ and the question arises why the initial energy density of the universe was so finely tuned to be equal to its critical value.

1.3 Magnetic Monopole Problem

This problem arises only if we combine the SBB Model with Grand Unified Theories³ (GUTs) of strong, weak and electromagnetic interactions. In accordance with GUTs, the universe underwent⁴ a phase transition during which the GUT gauge symmetry group, G , broke down to the Standard Model gauge group, G_S . This breaking was due to the fact that, at a critical temperature T_c , an appropriate higgs field, ϕ , developed a non-zero vacuum expectation value (vev). Assuming that this phase transition was a second order one, we have $\langle\phi\rangle(T) \approx \langle\phi\rangle(T=0)(1 - T^2/T_c^2)^{1/2}$, $m_H(T) \approx \lambda\langle\phi\rangle(T)$, for the temperature dependent vev and mass of the higgs field respectively at $T \leq T_c$ (λ is an appropriate higgs coupling constant).

The GUT phase transition produces magnetic monopoles⁵ which are localized deviations from the vacuum with radius $\sim M_X^{-1}$, energy $\sim M_X/\alpha_G$ and $\phi = 0$ at their center (M_X is the GUT mass scale and $\alpha_G = g_G^2/4\pi$ with g_G being the GUT gauge coupling constant). The vev of the higgs field on a sphere, S^2 , with radius $\gg M_X^{-1}$ around the monopole lies on the vacuum manifold G/G_S and we, thus, obtain a mapping: $S^2 \longrightarrow G/G_S$. If this mapping is homotopically non-trivial the topological stability of the magnetic monopole is guaranteed.

Monopoles can be produced when the fluctuations of ϕ over $\phi = 0$ between the vacua at $\pm\langle\phi\rangle(T)$ cease to be frequent. This happens when the free energy needed for ϕ to fluctuate from $\langle\phi\rangle(T)$ to zero in a region of radius equal to the higgs correlation length $\xi(T) = m_H^{-1}(T)$ exceeds T . This condition reads $(4\pi/3)\xi^3\Delta V \gtrsim T$, where $\Delta V \sim \lambda^2\langle\phi\rangle^4$ is the difference in free energy density between $\phi = 0$ and $\phi = \langle\phi\rangle(T)$. The Ginzburg temperature⁶, T_G , corresponds to the saturation of this inequality. So, at $T \lesssim T_G$, the fluctuations over $\phi = 0$ stop and $\langle\phi\rangle$ settles on the vacuum manifold G/G_S . At T_G , the universe splits into regions of size $\xi_G \sim (\lambda^2 T_c)^{-1}$, the higgs correlation length at T_G , with the higgs field more or less aligned in each region. Monopoles are produced at the corners where such regions meet (Kibble⁷ mechanism) and their number density is estimated to be $n_M \sim p\xi_G^{-3} \sim p\lambda^6 T_c^3$, where $p \sim 1/10$ is a geometric factor. The ‘relative’ monopole number density then turns out to be $r_M = n_M/T^3 \sim 10^{-6}$. We can derive a lower bound on r_M by pure causality considerations. The higgs field ϕ cannot be correlated at distances bigger than the particle horizon size, $2t_G$, at T_G . This gives the causality bound

$$n_M \gtrsim \frac{p}{\frac{4\pi}{3}(2t_G)^3} , \quad (4)$$

which implies that $r_M \gtrsim 10^{-10}$.

The subsequent evolution of monopoles, after T_G , is governed by the equation⁸ (overdots denote derivation with respect to cosmic time)

$$\frac{dn_M}{dt} = -Dn_M^2 - 3\frac{\dot{a}}{a}n_M, \quad (5)$$

where the first term in the right hand side (with D being an appropriate constant) describes the dilution of monopoles due to monopole-antimonopole annihilation while the second term corresponds to their dilution by the cosmological expansion. The monopole-antimonopole annihilation proceeds as follows. Monopoles diffuse towards antimonopoles through the plasma of charged particles, capture each other in Bohr orbits and eventually annihilate. The annihilation is effective provided the mean free path of the monopoles in the plasma of charged particles does not exceed their capture distance. This happens at cosmic temperatures $T \gtrsim 10^{12}$ GeV. The overall result is that, if the initial relative magnetic monopole density $r_{M,\text{in}} \gtrsim 10^{-9}$ ($\lesssim 10^{-9}$), the final one $r_{M,\text{fin}} \sim 10^{-9}$ ($\sim r_{M,\text{in}}$). This combined with the causality bound yields $r_{M,\text{fin}} \gtrsim 10^{-10}$. However, the requirement that monopoles do not dominate the energy density of the universe at nucleosynthesis gives

$$r_M(T \approx 1 \text{ MeV}) \lesssim 10^{-19}, \quad (6)$$

and we have a discrepancy of about ten orders of magnitude.

1.4 Density Fluctuations

For structure formation⁹ in the universe, we need a primordial density perturbation, $\delta\rho/\rho$, at all scales with a nearly flat spectrum¹⁰. We also need some explanation of the temperature fluctuations, $\delta T/T$, of CBR observed by COBE² at angles $\theta \gtrsim \theta_d \approx 2^\circ$ which violate causality (see Sec.1.1).

Let us expand $\delta\rho/\rho$ in plane waves

$$\frac{\delta\rho}{\rho}(\bar{x}, t) = \int d^3k \delta_{\bar{k}}(t) e^{i\bar{k}\bar{x}}, \quad (7)$$

where $\bar{x}$ is a comoving vector in 3-space and $\bar{k}$ is the comoving wave vector with $k = |\bar{k}|$ being the comoving wave number ($\lambda = 2\pi/k$ is the comoving wave length whereas the physical wave length is $\lambda_{\text{phys}} = a(t)\lambda$). For $\lambda_{\text{phys}} \leq H^{-1}$, the time evolution of $\delta_{\bar{k}}$ is described by the Newtonian equation

$$\ddot{\delta}_{\bar{k}} + 2H\dot{\delta}_{\bar{k}} + \frac{v_s^2 k^2}{a^2} \delta_{\bar{k}} = 4\pi G \rho \delta_{\bar{k}}, \quad (8)$$

where the second term in the left hand side comes from the cosmological expansion and the third is the ‘pressure’ term (v_s is the velocity of sound given by $v_s^2 = dp/d\rho$, where p is the mean pressure). The right hand side of this equation corresponds to the gravitational attraction.

For the moment, let us put $H=0$ (static universe). In this case, there exists a characteristic wave number k_J , the Jeans wave number, given by $k_J^2 = 4\pi G a^2 \rho / v_s^2$ and having the following property. For $k \geq k_J$, pressure dominates over gravitational attraction and the density perturbations just oscillate, whereas, for $k \leq k_J$, gravitational attraction dominates and the density perturbations grow exponentially. In particular, for $p=0$ (cold dark matter), $v_s = 0$ and all scales are Jeans unstable with

$$\delta_{\vec{k}} \propto \exp(t/\tau) , \quad \tau = (4\pi G \rho)^{-1/2} . \quad (9)$$

Now let us take $H \neq 0$. Since the cosmological expansion pulls the particles apart, we get a smaller growth:

$$\delta_{\vec{k}} \propto a(t) \propto t^{2/3} , \quad (10)$$

in the matter dominated case. For a radiation dominated universe ($p \neq 0$), we get essentially no growth of the density perturbations. This means that, in order to have structure formation in the universe, which requires $\delta\rho/\rho \sim 1$, we must have

$$\left(\frac{\delta\rho}{\rho}\right)_{\text{equ}} \sim 4 \times 10^{-5} (\Omega_0 h)^{-2} , \quad (11)$$

at the equidensity point (where the energy densities of matter and radiation coincide), since the available growth factor for perturbations is given by $a_0/a_{\text{equ}} \sim 2.5 \times 10^4 (\Omega_0 h)^2$. Here $\Omega_0 = \rho_0/\rho_c$, where ρ_0 is the present energy density of the universe. The question then is where these primordial density fluctuations originate from.

2 Inflation

Inflation^{11,12} is an idea which solves simultaneously all four cosmological puzzles and can be summarized as follows. Suppose there is a real scalar field ϕ (the inflaton) with (symmetric) potential energy density $V(\phi)$ which is quite ‘flat’ near $\phi = 0$ and has minima at $\phi = \pm\langle\phi\rangle$ with $V(\pm\langle\phi\rangle) = 0$. At high enough T ’s, $\phi = 0$ in the universe due to the temperature corrections in $V(\phi)$. As T drops, the effective potential density approaches the $T=0$ potential but a little potential barrier separating the local minimum at $\phi = 0$ and the vacua at $\phi = \pm\langle\phi\rangle$ still remains. At some point, ϕ tunnels out to $\phi_1 \ll \langle\phi\rangle$ and a

bubble with $\phi = \phi_1$ is created in the universe. The field then rolls over to the minimum of $V(\phi)$ very slowly (due to the flatness of the potential). During this slow roll-over, the energy density $\rho \approx V(\phi = 0) \equiv V_0$ remains essentially constant for quite some time. The Lagrangian density

$$L = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) \quad (12)$$

gives the energy-momentum tensor

$$T_\mu{}^\nu = -\partial_\mu \phi \partial^\nu \phi + \delta_\mu{}^\nu \left(\frac{1}{2} \partial_\lambda \phi \partial^\lambda \phi - V(\phi) \right), \quad (13)$$

which during the slow roll-over takes the form $T_\mu{}^\nu \approx -V_0 \delta_\mu{}^\nu$. This means that $\rho \approx -p \approx V_0$, i.e., the pressure p is negative and equal in magnitude with the energy density ρ , which is consistent with the continuity equation $\dot{\rho} = -3H(\rho + p)$. Since, as we will see, $a(t)$ grows very fast, the ‘curvature’ term, k/a^2 , in Eq.3 becomes subdominant and we get

$$H^2 \equiv \left(\frac{\dot{a}}{a} \right)^2 = \frac{8\pi G}{3} V_0, \quad (14)$$

which gives $a(t) \propto e^{Ht}$, $H^2 = (8\pi G/3)V_0 = \text{constant}$. So the bubble expands exponentially for some time and $a(t)$ grows by a factor

$$\frac{a(t_f)}{a(t_i)} = \exp H(t_f - t_i) \equiv \exp H\tau, \quad (15)$$

between an initial (t_i) and a final (t_f) cosmic time.

The inflationary scheme just described, which is known as the new¹³ inflationary scenario (with the inflaton field starting from the origin, $\phi=0$), is certainly not the only realization of the idea of inflation. Another interesting possibility is to consider the universe as it emerges at the Planck time $t_P = m_P^{-1}$, $m_P = M_P/\sqrt{8\pi}$ with $M_P = 1.22 \times 10^{19}$ GeV, where the fluctuations of gravity cease to exist. We can imagine a region of size $\ell_P \approx m_P^{-1}$ where the inflaton field acquires a large and almost uniform value and carries negligible kinetic energy. Under certain circumstances this region can inflate (exponentially expand) as ϕ rolls down towards its vacuum value. This type of inflation with the inflaton starting from large values is known as the chaotic¹⁴ inflationary scenario.

We will now show that, with an adequate number of e-foldings, $N = H\tau$, the first three cosmological puzzles are easily resolved (we will leave the question of density perturbations for later).

2.1 Resolution of the Horizon Problem

The particle horizon during inflation (exponential expansion)

$$d(t) = e^{Ht} \int_{t_i}^t \frac{dt'}{e^{Ht'}} \approx H^{-1} \exp H(t - t_i) , \quad (16)$$

for $t - t_i \gg H^{-1}$, grows as fast as $a(t)$. At the end of inflation ($t = t_f$), $d(t_f) \approx H^{-1} \exp H\tau$ and the field ϕ starts oscillating about the minimum of the potential at $\phi = \langle \phi \rangle$. It then decays and ‘reheats’¹⁵ the universe at a temperature $T_r \sim 10^9$ GeV (see Sec.10). The universe, after that, goes back to normal big bang cosmology. The horizon $d(t_f)$ is stretched during the period of ϕ -oscillations by some factor $\sim 10^9$ depending on details and between T_r and the present era by a factor T_r/T_0 . So it finally becomes equal to $H^{-1} e^{H\tau} 10^9 (T_r/T_0)$, which should exceed $2H_0^{-1}$ in order to solve the horizon problem. Taking $V_0 \approx M_X^4$, $M_X \sim 10^{16}$ GeV, we see that, with $N = H\tau \gtrsim 55$, the horizon problem is evaded.

2.2 Resolution of the Flatness Problem

The curvature term of the Friedmann equation, at present, is given by

$$\frac{k}{a^2} \approx \left(\frac{k}{a^2} \right)_{bi} e^{-2H\tau} 10^{-18} \left(\frac{10^{-13} \text{ GeV}}{10^9 \text{ GeV}} \right)^2 , \quad (17)$$

where the terms in the right hand side correspond to the curvature term before inflation, and its growth factors during inflation, during ϕ -oscillations and after ‘reheating’ respectively. Assuming $(k/a^2)_{bi} \sim (8\pi G/3)\rho \sim H^2$ ($\rho \approx V_0$), we get $k/a_0^2 H_0^2 \sim 10^{48} e^{-2H\tau}$ which gives $(\rho_0 - \rho_c)/\rho_c \equiv \Omega_0 - 1 = k/a_0^2 H_0^2 \ll 1$, for $H\tau \gg 55$. In fact, strong inflation implies that the present universe is flat with a great accuracy.

2.3 Resolution of the Monopole Problem

It is obvious that, with a number of e-foldings $\gtrsim 55$, the primordial monopole density is diluted by at least 70 orders of magnitude and they become totally irrelevant. Also, since $T_r \ll m_M$, there is no production of magnetic monopoles after ‘reheating’.

3 Detailed Analysis of Inflation

The Hubble parameter is not exactly constant during inflation as we, naively, assumed so far. It actually depends on the value of ϕ :

$$H^2(\phi) = \frac{8\pi G}{3} V(\phi) . \quad (18)$$

To find the evolution equation for ϕ during inflation, we vary the action

$$S = \int \sqrt{-\det(g)} d^4x \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) + M(\phi) \right) , \quad (19)$$

where g is the metric tensor and $M(\phi)$ represents the coupling of ϕ to ‘light’ matter causing its decay. We find

$$\ddot{\phi} + 3H\dot{\phi} + \Gamma_\phi \dot{\phi} + V'(\phi) = 0 , \quad (20)$$

where the prime denotes derivation with respect to ϕ and Γ_ϕ is the decay width¹⁶ of the inflaton. Assume, for the moment, that the decay time of ϕ , $t_d = \Gamma_\phi^{-1}$, is much greater than H^{-1} , the expansion time for inflation. Then the term $\Gamma_\phi \dot{\phi}$ can be ignored and Eq.20 reduces to

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0 . \quad (21)$$

Inflation is by definition the situation where $\ddot{\phi}$ is subdominant to the ‘friction’ term $3H\dot{\phi}$ in this equation (and the kinetic energy density is subdominant to the potential energy density). Eq.21 then further reduces to the inflationary equation¹⁷

$$3H\dot{\phi} = -V'(\phi) , \quad (22)$$

which gives

$$\ddot{\phi} = -\frac{V''(\phi)\dot{\phi}}{3H(\phi)} + \frac{V'(\phi)}{3H^2(\phi)} H'(\phi) \dot{\phi} . \quad (23)$$

Comparing the two terms in the right hand side of this equation with the ‘friction’ term in Eq.21, we obtain the conditions for inflation (slow roll conditions):

$$\eta \equiv \frac{M_P^2}{8\pi} \left| \frac{V''(\phi)}{V(\phi)} \right| \leq 1 , \quad \epsilon \equiv \frac{M_P^2}{16\pi} \left(\frac{V'(\phi)}{V(\phi)} \right)^2 \leq 1 . \quad (24)$$

The end of the slow ‘roll-over’ occurs when either of these inequalities is saturated. If ϕ_f is the value of ϕ at the end of inflation, then $t_f \sim H^{-1}(\phi_f)$.

The number of e-foldings during inflation can be calculated as follows:

$$N(\phi_i \rightarrow \phi_f) \equiv \ell n \left(\frac{a(t_f)}{a(t_i)} \right) = \int_{t_i}^{t_f} H dt = \int_{\phi_i}^{\phi_f} \frac{H(\phi)}{\dot{\phi}} d\phi = - \int_{\phi_i}^{\phi_f} \frac{3H^2(\phi) d\phi}{V'(\phi)}, \quad (25)$$

where Eqs.15, 22 and the definition of $H = \dot{a}/a$ were used. For simplicity, we can shift the field ϕ so that the global minimum of the potential is displaced at $\phi = 0$. Then, if $V(\phi) = \lambda\phi^\nu$ during inflation, we have

$$N(\phi_i \rightarrow \phi_f) = - \int_{\phi_i}^{\phi_f} \frac{3H^2(\phi) d\phi}{V'(\phi)} = -8\pi G \int_{\phi_i}^{\phi_f} \frac{V(\phi) d\phi}{V'(\phi)} = \frac{4\pi G}{\nu} (\phi_i^2 - \phi_f^2). \quad (26)$$

Assuming that $\phi_i \gg \phi_f$, this reduces to $N(\phi) = (4\pi G/\nu)\phi^2$.

4 Coherent Field Oscillations

After the end of inflation at cosmic time t_f , the term $\ddot{\phi}$ takes over and Eq.21 reduces to $\ddot{\phi} + V'(\phi) = 0$, which means that ϕ starts oscillating coherently about the global minimum of the potential. In reality, due to the ‘friction’ term, ϕ performs damped oscillations with a rate of energy density loss given by

$$\frac{d\rho}{dt} = \frac{d}{dt} \left(\frac{1}{2} \dot{\phi}^2 + V(\phi) \right) = -3H\dot{\phi}^2 = -3H(\rho + p), \quad (27)$$

where $\rho = \dot{\phi}^2/2 + V(\phi)$ and the pressure $p = \dot{\phi}^2/2 - V(\phi)$. Averaging p over one oscillation of ϕ , we write¹⁸ $\rho + p = \gamma\rho$. Eq.27 then becomes $\dot{\rho} = -3H\gamma\rho$, which gives $d\rho/\rho = -3\gamma da/a$ and $\rho \propto a^{-3\gamma}$. The Friedmann equation then takes the form $\dot{a}/a \propto a^{-3\gamma/2}$ and we obtain $a(t) \propto t^{2/3\gamma}$.

The number γ for an oscillating field can be written as (assuming a symmetric potential)

$$\gamma = \frac{\int_0^T \dot{\phi}^2 dt}{\int_0^T \rho dt} = \frac{\int_0^{\phi_{\max}} \dot{\phi} d\phi}{\int_0^{\phi_{\max}} (\rho/\dot{\phi}) d\phi}, \quad (28)$$

where T and $\phi_{\max}$ are the period and the amplitude of the oscillation respectively. From the equation $\rho = \dot{\phi}^2/2 + V(\phi) = V_{\max}$, where $V_{\max}$ is the maximal potential energy density, we obtain $\dot{\phi} = \sqrt{2(V_{\max} - V(\phi))}$. Substituting this in Eq.28 we get¹⁸

$$\gamma = \frac{2 \int_0^{\phi_{\max}} (1 - V/V_{\max})^{1/2} d\phi}{\int_0^{\phi_{\max}} (1 - V/V_{\max})^{-1/2} d\phi}. \quad (29)$$

For a potential of the simple form $V(\phi) = \lambda\phi^\nu$, γ is readily found to be given by $\gamma = 2\nu/(\nu+2)$. Consequently, in this case, $\rho \propto a^{-6\nu/(\nu+2)}$ and $a(t) \propto t^{(\nu+2)/3\nu}$. For $\nu = 2$, in particular, one has $\gamma=1$, $\rho \propto a^{-3}$, $a(t) \propto t^{2/3}$ and the oscillating field behaves like pressureless ‘matter’. This is not unexpected since a coherent oscillating massive free field corresponds to a distribution of static massive particles. For $\nu=4$, however, we obtain $\gamma = 4/3$, $\rho \propto a^{-4}$, $a(t) \propto t^{1/2}$ and the system resembles ‘radiation’. For $\nu = 6$, one has $\gamma = 3/2$, $\rho \propto a^{-4.5}$, $a(t) \propto t^{4/9}$ and the expansion is slower than in a ‘radiation’ dominated universe (the pressure is higher than in ‘radiation’).

5 Decay of the Field ϕ

Reintroducing the ‘decay’ term $\Gamma_\phi\dot{\phi}$, Eq.20 can be written as

$$\dot{\rho} = \frac{d}{dt} \left(\frac{1}{2}\dot{\phi}^2 + V(\phi) \right) = -(3H + \Gamma_\phi)\dot{\phi}^2, \quad (30)$$

which is solved^{15,18} by

$$\rho(t) = \rho_f \left(\frac{a(t)}{a(t_f)} \right)^{-3\gamma} \exp[-\gamma\Gamma_\phi(t - t_f)], \quad (31)$$

where ρ_f is the energy density at the end of inflation at cosmic time t_f . The second and third factors in the right hand side of this equation represent the dilution of the field energy due to the expansion of the universe and the decay of ϕ to light particles respectively.

All pre-existing ‘radiation’ (known as ‘old radiation’) was diluted by inflation, so the only ‘radiation’ present is the one produced by the decay of ϕ and is known as ‘new radiation’. Its energy density satisfies^{15,18} the equation

$$\dot{\rho}_r = -4H\rho_r + \gamma\Gamma_\phi\rho, \quad (32)$$

where the first term in the right hand side represents the dilution of radiation due to the cosmological expansion while the second one is the energy density transfer from ϕ to ‘radiation’. Taking $\rho_r(t_f)=0$, this equation gives^{15,18}

$$\rho_r(t) = \rho_f \left(\frac{a(t)}{a(t_f)} \right)^{-4} \int_{t_f}^t \left(\frac{a(t')}{a(t_f)} \right)^{4-3\gamma} e^{-\gamma\Gamma_\phi(t'-t_f)} \gamma\Gamma_\phi dt'. \quad (33)$$

For $t_f \ll t_d$ and $\nu = 2$, this expression is approximated by

$$\rho_r(t) = \rho_f \left(\frac{t}{t_f} \right)^{-8/3} \int_0^t \left(\frac{t'}{t_f} \right)^{2/3} e^{-\Gamma_\phi t'} dt', \quad (34)$$

which, using the formula

$$\int_0^u x^{p-1} e^{-x} dx = e^{-u} \sum_{k=0}^{\infty} \frac{u^{p+k}}{p(p+1) \cdots (p+k)} , \quad (35)$$

can be written as

$$\rho_r = \frac{3}{5} \rho \Gamma_{\phi} t \left[1 + \frac{3}{8} \Gamma_{\phi} t + \frac{9}{88} (\Gamma_{\phi} t)^2 + \cdots \right] , \quad (36)$$

with $\rho = \rho_f(t/t_f)^{-2} \exp(-\Gamma_{\phi} t)$ being the energy density of the field ϕ which performs damped oscillations and decays into ‘light’ particles.

The energy density of the ‘new radiation’ grows relative to the energy density of the oscillating field and becomes essentially equal to it at a cosmic time $t_d = \Gamma_{\phi}^{-1}$ as one can deduce from Eq.36. After this time, the universe enters into the radiation dominated era and the normal big bang cosmology is recovered. The temperature at t_d , $T_r(t_d)$, is historically called the ‘reheating’ temperature although no supercooling and subsequent reheating of the universe actually takes place. Using the time to temperature relation for a radiation dominated universe we find that

$$T_r = \left(\frac{45}{16\pi^3 g_*} \right)^{1/4} (\Gamma_{\phi} M_P)^{1/2} , \quad (37)$$

where g_* is the effective number of degrees of freedom. For a potential of the type $V(\phi) = \lambda \phi^{\nu}$, the total expansion of the universe during the period of damped field oscillations is

$$\frac{a(t_d)}{a(t_f)} = \left(\frac{t_d}{t_f} \right)^{\frac{\nu+2}{3\nu}} . \quad (38)$$

6 Density Perturbations

We are now ready to sketch how inflation solves the density fluctuation problem described in Sec.1.4. As a matter of fact, inflation not only homogenizes the universe but also provides us with the primordial density fluctuations necessary for the structure formation in the universe. To understand the origin of these fluctuations, we must first introduce the notion of ‘event horizon’. Our ‘event horizon’, at a cosmic time t , includes all points with which we will eventually communicate sending signals at t . The instantaneous (at cosmic time t) radius of the ‘event horizon’ is

$$d_e(t) = a(t) \int_t^{\infty} \frac{dt'}{a(t')} . \quad (39)$$

It is obvious, from this formula, that the ‘event horizon’ is infinite for matter or radiation dominated universe. For inflation, however, we obtain a slowly varying event horizon with radius $d_e(t) = H^{-1} < \infty$. Points, in our ‘event horizon’ at t , with which we can communicate sending signals at t , are eventually pulled away by the ‘exponential’ expansion and we cease to be able to communicate with them again emitting signals at later times. We say that these points (and the corresponding scales) crossed outside the event horizon. The situation is very similar to that of a black hole. Indeed, the exponentially expanding (de Sitter) space is like a black hole turned inside out. This means that we are inside and the black hole surrounds us from all sides. Then, exactly as in a black hole, there are quantum fluctuations of the ‘thermal type’ governed by the ‘Hawking temperature’^{19,20} $T_H = H/2\pi$. It turns out^{21,22} that the quantum fluctuations of all massless fields (the inflaton is nearly massless due to the ‘flatness’ of the potential) are $\delta\phi = H/2\pi = T_H$. These fluctuations of ϕ lead to energy density fluctuations $\delta\rho = V'(\phi)\delta\phi$. As the scale of this perturbations crosses outside the event horizon, they become²³ classical metric perturbations.

The evolution of these fluctuations outside the ‘inflationary horizon’ is quite subtle and involved due to the gauge freedom in general relativity. However, there is a simple gauge invariant quantity²⁴ $\zeta \approx \delta\rho/(\rho + p)$, which remains constant outside the horizon. Thus, the density fluctuation at any present physical (comoving) scale ℓ , $(\delta\rho/\rho)_\ell$, when this scale crosses inside the ‘post-inflationary’ particle horizon ($p=0$ at this instance) can be related to the value of ζ when the same scale crossed outside the inflationary event horizon (symbolically at $\ell \sim H^{-1}$). This latter value of ζ can be found using Eq.22 and turns out to be

$$\zeta|_{\ell \sim H^{-1}} = \left(\frac{\delta\rho}{\dot{\phi}^2} \right)_{\ell \sim H^{-1}} = \left(\frac{V'(\phi)H(\phi)}{2\pi\dot{\phi}^2} \right)_{\ell \sim H^{-1}} = - \left(\frac{9H^3(\phi)}{2\pi V'(\phi)} \right)_{\ell \sim H^{-1}} . \quad (40)$$

Taking into account an extra $2/5$ factor from the fact that the universe is matter dominated when the scale ℓ re-enters the horizon, we obtain

$$\left(\frac{\delta\rho}{\rho} \right)_\ell = \frac{16\sqrt{6}\pi}{5} \frac{V^{3/2}(\phi_\ell)}{M_P^3 V'(\phi_\ell)} . \quad (41)$$

The calculation of ϕ_ℓ , the value of the inflaton field when the ‘comoving’ scale ℓ crossed outside the event horizon, goes as follows. A ‘comoving’ (present physical) scale ℓ , at T_r , was equal to $\ell(a(t_d)/a(t_0)) = \ell(T_0/T_r)$. Its magnitude at the end of inflation ($t = t_f$) was $\ell(T_0/T_r)(a(t_f)/a(t_d)) = \ell(T_0/T_r)(t_f/t_d)^{(\nu+2)/3\nu} \equiv \ell_{\text{phys}}(t_f)$, where the potential $V(\phi) = \lambda\phi^\nu$ was assumed. The scale ℓ , when it crossed outside the inflationary horizon, was equal

to $H^{-1}(\phi_\ell)$. We, thus, obtain

$$H^{-1}(\phi_\ell)e^{N(\phi_\ell)} = \ell_{\text{phys}}(t_f) . \quad (42)$$

Solving this equation, one can calculate ϕ_ℓ and, subsequently, $N(\phi_\ell) \equiv N_\ell$, the number of e-foldings the scale ℓ suffered during inflation. In particular, for our present horizon scale $\ell \approx 2H_0^{-1} \sim 10^4$ Mpc (H_0 is the present value of the Hubble parameter), it turns out that $N_{H_0} \approx 50 - 60$.

Now, taking the potential $V(\phi) = \lambda\phi^4$, Eqs.26, 41 and 42 give

$$\left(\frac{\delta\rho}{\rho}\right)_\ell = \frac{4\sqrt{6}\pi}{5}\lambda^{1/2}\left(\frac{\phi_\ell}{M_P}\right)^3 = \frac{4\sqrt{6}\pi}{5}\lambda^{1/2}\left(\frac{N_\ell}{\pi}\right)^{3/2} . \quad (43)$$

The measurements of COBE ², $(\delta\rho/\rho)_{H_0} \approx 6 \times 10^{-5}$, then imply that $\lambda \approx 6 \times 10^{-14}$ for $N_{H_0} \approx 55$. Thus, we see that the inflaton must be a very weakly coupled field. In non-supersymmetric GUTs, the inflaton must necessarily be a gauge singlet field since otherwise radiative corrections will certainly make it strongly coupled. This is, undoubtedly, not a very satisfactory situation since we are forced to introduce an otherwise unmotivated extra *ad hoc* very weakly coupled gauge singlet. In supersymmetric GUTs, however, the inflaton could be identified ²⁵ with a conjugate pair of gauge non-singlet fields $\phi, \bar{\phi}$, already existing in the theory and causing the gauge symmetry breaking. Absence of strong radiative corrections from gauge interactions is guaranteed, in this case, by the mutual cancellation of the D-terms of these fields.

The spectrum of the density fluctuations emerging from inflation can also be analyzed. For definiteness, we will again take the potential $V(\phi) = \lambda\phi^\nu$. One then finds that $(\delta\rho/\rho)_\ell$ is proportional to $\phi_\ell^{(\nu+2)/2}$ which, combined with the fact that $N(\phi_\ell)$ is proportional to ϕ_ℓ^2 (see Eq.26), gives

$$\left(\frac{\delta\rho}{\rho}\right)_\ell = \left(\frac{\delta\rho}{\rho}\right)_{H_0} \left(\frac{N_\ell}{N_{H_0}}\right)^{\frac{\nu+2}{4}} . \quad (44)$$

The scale ℓ divided by the size of our present horizon ($\approx 10^4$ Mpc) should equal $\exp(N_\ell - N_{H_0})$. This gives $N_\ell/N_{H_0} = 1 + \ell n(\ell/10^4)^{1/N_{H_0}}$ which expanded around $\ell \approx 10^4$ Mpc and substituted in Eq.44 yields

$$\left(\frac{\delta\rho}{\rho}\right)_\ell = \left(\frac{\delta\rho}{\rho}\right)_{H_0} \left(\frac{\ell}{10^4 \text{ Mpc}}\right)^{\alpha_s} , \quad (45)$$

with $\alpha_s = (\nu + 2)/4N_{H_0}$. For $\nu = 4$, $\alpha_s \approx 0.03$ and the fluctuations are essentially scale independent.

7 Density Fluctuations in ‘Matter’

We will now discuss the evolution of the primordial density fluctuations after their scale enters the post-inflationary horizon. To this end, we introduce²⁶ the ‘conformal’ time, η , so that the Robertson-Walker metric takes the form of a conformally expanding Minkowski space:

$$ds^2 = -dt^2 + a^2(t) d\bar{r}^2 = a^2(\eta) (-d\eta^2 + d\bar{r}^2) , \quad (46)$$

where $\bar{r}$ is a comoving 3-vector. The Hubble parameter now takes the form $H = \dot{a}(t)/a(t) = a'(\eta)/a^2(\eta)$ and the Friedmann equation can be rewritten as

$$\frac{1}{a^2} \left(\frac{a'}{a} \right)^2 = \frac{8\pi G}{3} \rho , \quad (47)$$

where primes denote derivation with respect to the ‘conformal’ time η . The continuity equation takes the form $\rho' = -3\tilde{H}(\rho + p)$ with $\tilde{H} = a'/a$. For a matter dominated universe, $\rho \propto a^{-3}$ which gives $a = (\eta/\eta_0)^2$ and $a'/a = 2/\eta$ (η_0 is the present value of η).

The ‘Newtonian’ Eq.8 can now be written in the form

$$\delta''_{\bar{k}}(\eta) + \frac{a'}{a} \delta'_{\bar{k}}(\eta) - 4\pi G \rho a^2 \delta_{\bar{k}}(\eta) = 0 , \quad (48)$$

and the growing (Jeans unstable) mode $\delta_{\bar{k}}(\eta)$ is proportional to η^2 and can be expressed²⁷ as

$$\delta_{\bar{k}}(\eta) = \epsilon_H \left(\frac{k\eta}{2} \right)^2 \hat{s}(\bar{k}) , \quad (49)$$

where $\hat{s}(\bar{k})$ is a Gaussian random variable satisfying

$$\langle \hat{s}(\bar{k}) \rangle = 0 , \quad \langle \hat{s}(\bar{k}) \hat{s}(\bar{k}') \rangle = \frac{1}{k^3} \delta(\bar{k} - \bar{k}') , \quad (50)$$

and ϵ_H is the amplitude of the perturbation when its scale crosses inside the ‘post-inflationary horizon’. The latter can be seen as follows. A ‘comoving’ (present physical) length ℓ crosses inside the post-inflationary horizon when $a\ell/2\pi = H^{-1} = a^2/a'$ which gives $\ell/2\pi \equiv k^{-1} = a/a' = \eta_H/2$ or $k\eta_H/2 = 1$, where η_H is the conformal time at horizon crossing. This means that, at horizon crossing, $\delta_{\bar{k}}(\eta_H) = \epsilon_H \hat{s}(\bar{k})$. For scale invariant perturbations, the amplitude ϵ_H is constant. The gauge invariant perturbations of the ‘scalar gravitational potential’ are given²⁶ by the Poisson’s equation,

$$\Phi = -4\pi G \frac{a^2}{k^2} \rho \delta_{\bar{k}}(\eta) . \quad (51)$$

From the Friedmann Eq.47, we then obtain

$$\Phi = -\frac{3}{2}\epsilon_H \hat{s}(\bar{k}) . \quad (52)$$

The spectrum of the density perturbations can be characterized by the correlation function

$$\xi(\bar{r}) \equiv \langle \tilde{\delta}^*(\bar{x}, \eta) \tilde{\delta}(\bar{x} + \bar{r}, \eta) \rangle , \quad (53)$$

where

$$\tilde{\delta}(\bar{x}, \eta) = \int d^3k \delta_{\bar{k}}(\eta) e^{i\bar{k}\bar{x}} . \quad (54)$$

Substituting Eq.49 in Eq.53 and using Eq.50, we obtain

$$\xi(\bar{r}) = \int d^3k e^{-i\bar{k}\bar{r}} \epsilon_H^2 \left(\frac{k\eta}{2} \right)^4 \frac{1}{k^3} , \quad (55)$$

and, thus, the spectral function $P(k, \eta) = \epsilon_H^2 (\eta^4/16)k$ is proportional to k for ϵ_H constant. We say that, in this case, the ‘spectral index’ $n = 1$ and we have a Harrison-Zeldovich¹⁰ flat spectrum. In the general case, $P \propto k^n$ with $n = 1 - 2\alpha_s$ (see Eq.45). For $V(\phi) = \lambda\phi^4$, we get $n \approx 0.94$.

8 Temperature Fluctuations

The density inhomogeneities produce temperature fluctuations in the CBR. For angles $\theta \gtrsim 2^\circ$, the dominant effect is the scalar Sachs-Wolfe²⁸ effect. Density perturbations on the ‘last scattering surface’ cause scalar gravitational potential fluctuations, Φ , which, in turn, produce temperature fluctuations in the CBR. The physical reason is that regions with a deep gravitational potential will cause the photons to lose energy as they climb up the well and, thus, appear cooler. For $\theta \lesssim 2^\circ$, the dominant effects are: i) Motion of the last scattering surface causing Doppler shifts, and ii) Intrinsic fluctuations of the photon temperature, T_γ , which are more difficult to calculate since they depend on microphysics, the ionization history, photon streaming and other effects.

The temperature fluctuations at an angle θ due to the scalar Sachs-Wolfe effect turn out²⁸ to be given by $(\delta T/T)_\theta = -\Phi_\ell/3$, where ℓ is the ‘comoving’ scale on the ‘last scattering surface’ which subtends the angle θ [$\ell \approx 100 h^{-1}(\theta/\text{degrees}) \text{ Mpc}$] and Φ_ℓ the corresponding scalar gravitational potential fluctuations. From Eq.52, we then obtain $(\delta T/T)_\theta = (\epsilon_H/2)\hat{s}(\bar{k})$, which

using Eq.49 becomes

$$\left(\frac{\delta T}{T}\right)_\theta = \frac{1}{2}\delta_k(\eta_H) = \frac{1}{2}\left(\frac{\delta\rho}{\rho}\right)_{\ell \sim 2\pi k^{-1}}. \quad (56)$$

The COBE scale (our present horizon) corresponds to $\theta \approx 60^\circ$. Eqs.26, 41 and 56 give

$$\left(\frac{\delta T}{T}\right)_\ell \propto \left(\frac{\delta\rho}{\rho}\right)_\ell \propto \frac{V^{3/2}(\phi_\ell)}{M_P^3 V'(\phi_\ell)} \propto N_\ell^{\frac{\nu+2}{4}}. \quad (57)$$

Analyzing the temperature fluctuations in spherical harmonics we can obtain the quadrupole anisotropy due to the scalar Sachs-Wolfe effect,

$$\left(\frac{\delta T}{T}\right)_{Q-S} = \left(\frac{32\pi}{45}\right)^{1/2} \frac{V^{3/2}(\phi_\ell)}{M_P^3 V'(\phi_\ell)}. \quad (58)$$

For $V(\phi) = \lambda\phi^\nu$, this becomes

$$\left(\frac{\delta T}{T}\right)_{Q-S} = \left(\frac{32\pi}{45}\right)^{1/2} \frac{\lambda^{1/2} \phi_\ell^{\frac{\nu+2}{2}}}{\nu M_P^3} = \left(\frac{32\pi}{45}\right)^{1/2} \frac{\lambda^{1/2}}{\nu M_P^3} \left(\frac{\nu M_P^2}{4\pi}\right)^{\frac{\nu+2}{4}} N_\ell^{\frac{\nu+2}{4}}. \quad (59)$$

Comparing this with the COBE² measurements, $(\delta T/T)_Q \approx 6.6 \times 10^{-6}$, we obtain $\lambda \approx 6 \times 10^{-14}$, for $\nu = 4$ and number of e-foldings suffered by our present horizon scale during inflation $N_{\ell \sim H_0^{-1}} \equiv N_Q \approx 55$.

There are also ‘tensor’²⁹ (gravitational wave) fluctuations in the temperature of CBR. The quadrupole tensor anisotropy is

$$\left(\frac{\delta T}{T}\right)_{Q-T} \approx 0.77 \frac{V^{1/2}(\phi_\ell)}{M_P^2}. \quad (60)$$

The total quadrupole anisotropy is given by

$$\left(\frac{\delta T}{T}\right)_Q = \left[\left(\frac{\delta T}{T}\right)_{Q-S}^2 + \left(\frac{\delta T}{T}\right)_{Q-T}^2 \right]^{1/2}, \quad (61)$$

and the ratio

$$r = \frac{(\delta T/T)_{Q-T}^2}{(\delta T/T)_{Q-S}^2} \approx 0.27 \left(\frac{M_P V'(\phi_\ell)}{V(\phi_\ell)} \right)^2. \quad (62)$$

For $V(\phi) = \lambda\phi^\nu$, we obtain $r \approx 3.4 \nu/N_H \ll 1$, and the ‘tensor’ contribution to the temperature fluctuations of the CBR is negligible.

9 Hybrid Inflation

9.1 The non Supersymmetric Version

The most important disadvantage of the inflationary scenarios described so far is that they need extremely small coupling constants in order to reproduce the results of COBE². This difficulty was overcome some years ago by Linde³⁰ who proposed, in the context of non-supersymmetric GUTs, a clever inflationary scenario known as hybrid inflation. The idea was to use two real scalar fields χ and σ instead of one that was normally used. The field χ provides the vacuum energy which drives inflation while σ is the slowly varying field during inflation. The main advantage of this scenario is that it can reproduce the observed temperature fluctuations of the CBR with ‘natural’ values of the parameters in contrast to previous realizations of inflation (like the ‘new’¹³ or ‘chaotic’¹⁴ inflationary scenarios). The potential utilized by Linde is

$$V(\chi, \sigma) = \kappa^2 \left(M^2 - \frac{\chi^2}{4} \right)^2 + \frac{\lambda^2 \chi^2 \sigma^2}{4} + \frac{m^2 \sigma^2}{2} , \quad (63)$$

where κ , λ are dimensionless positive coupling constants and M , m mass parameters. The vacua lie at $\langle \chi \rangle = \pm 2M$, $\langle \sigma \rangle = 0$. Putting $m=0$, for the moment, we observe that the potential possesses an exactly flat direction at $\chi = 0$ with $V(\chi = 0, \sigma) = \kappa^2 M^4$. The mass squared of the field χ along this flat direction is given by $m_\chi^2 = -\kappa^2 M^2 + \frac{1}{2} \lambda^2 \sigma^2$ and remains non-negative for $\sigma \geq \sigma_c = \sqrt{2} \kappa M / \lambda$. This means that, at $\chi = 0$ and $\sigma \geq \sigma_c$, we obtain a valley of minima with flat bottom. Reintroducing the mass parameter m in Eq.63, we observe that this valley acquires a non-zero slope. A region of the universe, where χ and σ happen to be almost uniform with negligible kinetic energies and with values close to the bottom of the valley of minima, follows this valley in its subsequent evolution and undergoes inflation. The quadrupole anisotropy of CBR produced during this inflation can be estimated, from Eq.58, to be

$$\left(\frac{\delta T}{T} \right)_Q \approx \left(\frac{16\pi}{45} \right)^{1/2} \frac{\lambda \kappa^2 M^5}{M_P^3 m^2} . \quad (64)$$

The COBE² result, $(\delta T/T)_Q \approx 6.6 \times 10^{-6}$, can then be reproduced with $M \approx 2.86 \times 10^{16}$ GeV (the supersymmetric GUT vev) and $m \approx 1.3 \kappa \sqrt{\lambda} \times 10^{15}$ GeV $\sim 10^{12}$ GeV for $\kappa, \lambda \sim 10^{-2}$. Inflation terminates abruptly at $\sigma = \sigma_c$ and is followed by a ‘waterfall’, i.e., a sudden entrance into an oscillatory phase about a global minimum. Since the system can fall into either of the two available global minima with equal probability, topological defects are

copiously produced if they are predicted by the particular particle physics model one is considering.

9.2 The Supersymmetric Version

The hybrid inflationary scenario is ³¹ ‘tailor made’ for application to supersymmetric GUTs except that the mass of σ , m , is unacceptably large for supersymmetry, where all scalar fields acquire masses of order $m_S \sim 1$ TeV from soft supersymmetry breaking. To see this, consider a supersymmetric GUT with a (semi-simple) gauge group G of rank ≥ 5 with $G \rightarrow G_S$ (the Standard Model gauge group) at a scale $M \sim 10^{16}$ GeV. The spectrum of the theory below M is assumed to coincide with the Minimal Supersymmetric Standard Model (MSSM) spectrum plus Standard Model singlets so that the successful predictions for α_s , $\sin^2\theta_W$ are retained. The theory may also possess global symmetries. The breaking of G is achieved through the superpotential

$$W = \kappa S(-M^2 + \bar{\phi}\phi), \quad (65)$$

where $\bar{\phi}, \phi$ is a conjugate pair of Standard Model singlet left handed superfields which belong to non-trivial representations of G and reduce its rank by their vevs and S is a gauge singlet left handed superfield. The coupling constant κ and the mass parameter M can be made real and positive by suitable redefinitions of the phases of the superfields. This superpotential is the most general renormalizable superpotential consistent with a $U(1)$ R symmetry under which $W \rightarrow e^{i\theta} W$, $S \rightarrow e^{i\theta} S$, $\bar{\phi}\phi \rightarrow \bar{\phi}\phi$ and gives the potential

$$V = \kappa^2 |M^2 - \bar{\phi}\phi|^2 + \kappa^2 |S|^2 (|\phi|^2 + |\bar{\phi}|^2) + \text{D - terms}. \quad (66)$$

Restricting ourselves to the D-flat direction $\bar{\phi}^* = \phi$ which contains the supersymmetric minima and performing appropriate gauge and R transformations, we can bring S , $\bar{\phi}$, ϕ on the real axis, i.e., $S \equiv \sigma/\sqrt{2}$, $\bar{\phi} = \phi \equiv \chi/2$, where σ , χ are normalized real scalar fields. The potential then takes the form in Eq.63 with $\kappa = \lambda$ and $m = 0$ and, thus, Linde’s potential for hybrid inflation is almost obtainable from supersymmetric GUTs but without the mass term of σ which is, however, of crucial importance since it provides the slope of the valley of minima necessary for inflation.

One way to obtain a valley of minima useful for inflation is ³² to replace the renormalizable trilinear term in the superpotential W in Eq.65 by the next order non-renormalizable coupling. Another way, which we will adopt here, is ³³ to keep the renormalizable superpotential in Eq.65 and use the radiative

corrections along the inflationary valley ($\phi = \bar{\phi} = 0$, $S > S_c \equiv M$). In fact, due to supersymmetry breaking by the ‘vacuum’ energy density, $\kappa^2 M^4$, along this valley, there are important radiative corrections. At one loop, and for S sufficiently larger than S_c , the inflationary potential is given^{33,34} by

$$V_{\text{eff}}(S) = \kappa^2 M^4 \left[1 + \frac{\kappa^2}{16\pi^2} \left(\ell n \left(\frac{\kappa^2 S^2}{\Lambda^2} \right) + \frac{3}{2} - \frac{S_c^4}{12S^4} + \dots \right) \right], \quad (67)$$

where Λ is a suitable mass renormalization scale. Using this effective potential and Eq.58, one finds that the cosmic microwave quadrupole anisotropy $(\delta T/T)_Q \approx 8\pi(N_Q/45)^{1/2}(x_Q/y_Q)(M/M_P)^2$. Here $y_Q = x_Q(1 - 7/(12x_Q^2) + \dots)$ with $x_Q = S_Q/M$, and S_Q is the value of the scalar field S when the scale which evolved to the present horizon size crossed outside the de Sitter (inflationary) horizon. Also from Eq.67, one finds $\kappa \approx (8\pi^{3/2}/\sqrt{N_Q}) y_Q (M/M_P)$.

The inflationary phase ends as S approaches S_c from above. Writing $S = xS_c$, $x = 1$ corresponds to the phase transition from G to G_S which, as it turns out, more or less coincides with the end of the inflationary phase (this is checked by noting the amplitude of the quantities ϵ and η in Eq.24). Indeed, the 50–60 e-foldings needed for the inflationary scenario can be realized even with small values of x_Q . For definiteness, we will take $x_Q \approx 2$ (see Sec.12). From COBE² one then obtains $M \approx 5.5 \times 10^{15} \text{ GeV}$ and $\kappa \approx 4.5 \times 10^{-3}$ for $N_Q \approx 56$. Moreover, the primordial density fluctuation spectral index $n \simeq 0.98$. We see that the relevant part of inflation takes place at $S \sim 10^{16} \text{ GeV}$. An important consequence of this is^{31,34,35} that the supergravity corrections can be negligible.

In conclusion, it is important to note that the superpotential W in Eq.65 leads to hybrid inflation in a ‘natural’ way. This means that a) there is no need of very small coupling constants, b) W is the most general renormalizable superpotential allowed by the gauge and R symmetries, and c) supersymmetry guarantees that the radiative corrections do not invalidate inflation. They rather provide a slope along the inflationary trajectory which drives the inflaton towards the supersymmetric vacua.

10 ‘Reheating’ in Supersymmetric Hybrid Inflation

In order to discuss the ‘reheating’ of the universe after inflation, we need to be a little more specific about the underlying particle physics model. To this end, we consider³⁴ a supersymmetric model based on the left-right symmetric gauge group $G_{LR} = SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$. Of course, it is anticipated that G_{LR} is embedded in a grand unified theory such as $SO(10)$ or $SU(3)_c \times SU(3)_L \times SU(3)_R$. The breaking of G_{LR} to G_S is achieved by the renormalizable superpotential in Eq.65, where $\phi, \bar{\phi}$ are now identified with the

Standard Model singlet components of a conjugate pair of $SU(2)_R \times U(1)_{B-L}$ doublet left handed superfields.

The inflaton (oscillating system) consists of the two complex scalar fields S and $\theta = (\delta\phi + \delta\bar{\phi})/\sqrt{2}$, where $\delta\phi = \phi - M$, $\delta\bar{\phi} = \bar{\phi} - M$, with mass $m_{infl} = \sqrt{2}\kappa M$. We mainly concentrate on the decay of θ . Its relevant coupling to ‘matter’ is provided³⁴ by the non-renormalizable superpotential coupling (in symbolic form)

$$\frac{1}{2} \left(\frac{M_{\nu^c}}{M^2} \right) \bar{\phi} \bar{\phi} \nu^c \nu^c, \quad (68)$$

where M_{ν^c} denotes the Majorana mass of the relevant right handed neutrino ν^c . Without loss of generality, we assume that the Majorana mass matrix of the right handed neutrinos has been brought to diagonal form with positive entries. Clearly, θ decays predominantly into the heaviest right handed neutrino permitted by phase space, i.e., with $M_{\nu^c} \leq m_{infl}/2$.

The field S is not important for reheating since it can rapidly decay³⁴ into higgsinos through the renormalizable superpotential term $\xi S h^{(1)} h^{(2)}$ allowed by gauge symmetry, where $h^{(1)}$, $h^{(2)}$ denote the electroweak higgs doublets which couple to the up and down type quarks respectively, and ξ is a suitable coupling constant. Note that, after supersymmetry breaking, S acquires^{34,36} an expectation value $\langle S \rangle \sim m_S$ ($m_S \sim 1$ TeV being the magnitude of supersymmetry breaking in the visible sector) and generates the μ term.

Following standard procedures (see Eq. 37), and assuming the MSSM spectrum, the ‘reheating’ temperature T_r is found to be given by

$$T_r \approx \frac{1}{7} (\Gamma_\theta M_P)^{1/2}, \quad (69)$$

where $\Gamma_\theta \approx (1/16\pi)(\sqrt{2}M_{\nu^c}/M)^2 \sqrt{2}\kappa M$ is the decay rate of θ . Substituting κ as a function of N_Q , y_Q and M , we find³⁴

$$T_r \approx \frac{1}{12} \left(\frac{56}{N_Q} \right)^{1/4} \sqrt{y_Q} M_{\nu^c}. \quad (70)$$

For $x_Q \approx 2$ and $N_Q \approx 56$, we have $T_r \approx M_{\nu^c}/9.23$. We will assume that T_r is restricted by the gravitino constraint³⁷, $T_r \lesssim 10^9$ GeV. Note that T_r is closely linked to the mass of the heaviest ν^c satisfying $M_{\nu^c} \leq m_{infl}/2$.

11 Baryogenesis via Leptogenesis

11.1 Primordial Leptogenesis

In the hybrid inflationary models under consideration here, it is not convenient to produce the baryon asymmetry of the universe (BAU) in the usual

way, i.e., through the decay of color 3, $\bar{3}$ fields (g, g^c). Some of the reasons are the following: i) For theories where leptons and quarks belong to different representations of the unifying gauge group G (which is the case, for example, for $G = G_{LR}$ or $SU(3)_c \times SU(3)_L \times SU(3)_R$), the baryon number can be made almost exactly conserved by imposing an appropriate discrete symmetry. In particular, for $G = G_{LR}$, we can impose³⁸ a discrete symmetry under which $q \rightarrow -q, q^c \rightarrow -q^c, \bar{q} \rightarrow -\bar{q}, \bar{q}^c \rightarrow -\bar{q}^c$ and all other superfields remain invariant ($q, q^c, \bar{q}, \bar{q}^c$ are superfields with the quantum numbers of the quarks, antiquarks and their conjugates respectively). ii) For theories where such a discrete symmetry is absent, we could, in principle, use as inflaton a pair of conjugate Standard Model singlet superfields $N, \bar{N}$ which decay into g, g^c . For $G = SU(3)_c \times SU(3)_L \times SU(3)_R$, for example, N ($\bar{N}$) could be the Standard Model singlet component of the $(1, \bar{3}, 3)$ ($(1, 3, \bar{3})$) superfields with zero $U(1)_{B-L}$ charge. But this is again unacceptable since the breaking of $SU(3)_c \times SU(3)_L \times SU(3)_R$ by the vevs of $N, \bar{N}$ predicts³⁹ magnetic monopoles which can then be copiously produced after inflation. Also $T_r \lesssim 10^9$ GeV (gravitino constraint³⁷) implies $m_g \lesssim 10^{10}$ GeV (from the coupling $(m_g/\langle N \rangle) N g g^c$) which leads to strong deviation from MSSM and possible proton decay problems.

So it is preferable to produce first a primordial lepton asymmetry⁴⁰ which can then be partially turned into the observed baryon asymmetry of the universe by the non-perturbative ‘sphaleron’ effects⁴¹ of the electroweak sector. In the particular model based on G_{LR} which we consider here, this is the only way to produce the BAU since the inflaton decays into right handed neutrinos. Their subsequent decay into ordinary higgs particles (higgsinos) and light leptons (sleptons) can produce the primordial lepton asymmetry. It is important, though, to ensure that this primordial lepton asymmetry is not erased by lepton number violating $2 \rightarrow 2$ scatterings such as $ll \rightarrow h^{(1)*} h^{(1)*}$ or $lh^{(1)} \rightarrow \bar{l} h^{(1)*}$ (l represents a lepton doublet) at all temperatures between T_r and 100 GeV. This is automatically satisfied since the lepton asymmetry is protected⁴² by supersymmetry at temperatures between T_r and $T \sim 10^7$ GeV, and for $T \lesssim 10^7$ GeV, these $2 \rightarrow 2$ scatterings are⁴³ well out of equilibrium.

The lepton asymmetry produced by the out-of-equilibrium decay ($M_{\nu_i^c} \gg T_r$) of the right handed neutrinos ν_i^c , which emerged from the inflaton decay, is calculated to be^{40,44}

$$\frac{n_L}{s} \approx -\frac{3}{16\pi} \frac{T_r}{m_{infl}} \sum_{l \neq i} g(r_{li}) \frac{\text{Im}(U M^{D'} M^{D'\dagger} U^\dagger)_{il}^2}{|\langle h^{(1)} \rangle|^2 (U M^{D'} M^{D'\dagger} U^\dagger)_{ii}}, \quad (71)$$

where n_L and s are the lepton number and entropy densities, $M^{D'}$ is the diagonal ‘Dirac’ mass matrix, U a unitary transformation so that $UM^{D'}$ is

the ‘Dirac’ mass matrix in the basis where the ‘Majorana’ mass matrix of ν^c ’s is diagonal (see Sec.12) and $|\langle h^{(1)} \rangle| \approx 174$ GeV for large $\tan\beta$. The function

$$g(r_{li}) = r_{li} \ell n \left(\frac{1 + r_{li}^2}{r_{li}^2} \right), \quad r_{li} = \frac{M_l}{M_i}, \quad (72)$$

with $g(r) \sim 1/r$ as $r \rightarrow \infty$. Here we took into account the following prefactors: i) At ‘reheating’, we have $n_{infl} m_{infl} = (\pi^2/30) g_* T_r^4$ (n_{infl} is the number density of inflatons) which together with the relation $s = (2\pi^2/45) g_* T_r^3$ implies that $n_{infl}/s = (3/4)(T_r/m_{infl})$. ii) Since each inflaton decays into two ν^c ’s, their number density $n_{\nu^c} = 2n_{infl}$ which then gives $n_{\nu^c}/s = (3/2)(T_r/m_{infl})$. iii) Supersymmetry gives an extra factor of two.

11.2 Sphaleron Effects

To see how the primordial lepton asymmetry partially turns into the observed BAU, we must first discuss the non-perturbative baryon (B -) and lepton (L -) number violation⁴⁵ in the Standard Model. Consider the electroweak gauge symmetry $SU(2)_L \times U(1)_Y$ in the limit where the Weinberg angle $\theta_W = 0$ and concentrate on $SU(2)_L$ (inclusion of $\theta_W \neq 0$ does not alter the conclusions). Also, for the moment, ignore the fermions and higgs fields so as to have a pure $SU(2)_L$ gauge theory. This theory has⁴⁶ infinitely many classical vacua which are topologically distinct and are characterized by a ‘winding number’ $n \in Z$. In the ‘temporal gauge’ ($A_0 = 0$), the remaining gauge freedom consists of time independent transformations and the vacuum corresponds to a pure gauge

$$A_i = \frac{i}{g} \partial_i g(\bar{x}) g^{-1}(\bar{x}), \quad (73)$$

where g is the $SU(2)_L$ gauge coupling constant, $\bar{x}$ belongs to 3-space, $i=1,2,3$, $g(\bar{x}) \in SU(2)_L$, and $g(\bar{x}) \rightarrow 1$ as $|\bar{x}| \rightarrow \infty$. Thus, the 3-space compactifies to a sphere S^3 and $g(\bar{x})$ defines a map: $S^3 \rightarrow SU(2)_L$ (with the $SU(2)_L$ group being topologically equivalent to S^3). These maps are classified into homotopy classes constituting the third homotopy group of S^3 , $\pi_3(S^3)$, and are characterized by a ‘winding number’

$$n = \int d^3x \epsilon^{ijk} \text{tr} (\partial_i g(\bar{x}) g^{-1}(\bar{x}) \partial_j g(\bar{x}) g^{-1}(\bar{x}) \partial_k g(\bar{x}) g^{-1}(\bar{x})). \quad (74)$$

The corresponding vacua are denoted as $|n\rangle$, $n \in Z$.

The tunneling amplitude from the vacuum $|n_-\rangle$ at $t = -\infty$ to the vacuum $|n_+\rangle$ at $t = +\infty$ is given by the functional integral

$$\langle n_+ | n_- \rangle = \int (dA) e^{-S(A)} \quad (75)$$

over all gauge field configurations satisfying the appropriate boundary conditions at $t = \pm\infty$. Performing a Wick rotation, $x_0 \equiv t \rightarrow -ix_4$, we can go to Euclidean space-time. Any Euclidean field configuration with finite action is characterized by an integer topological number known as the Pontryagin number

$$q = \frac{g^2}{16\pi^2} \int d^4x \operatorname{tr} \left(F^{\mu\nu} \tilde{F}_{\mu\nu} \right) , \quad (76)$$

with $\mu, \nu=1,2,3,4$ and $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\lambda\rho} F^{\lambda\rho}$ being the dual field strength. But $\operatorname{tr}(F^{\mu\nu} \tilde{F}_{\mu\nu}) = \partial^\mu J_\mu$, where J_μ is the ‘Chern-Simons current’ given by

$$J_\mu = \epsilon_{\mu\nu\alpha\beta} \operatorname{tr} \left(A^\nu F^{\alpha\beta} - \frac{2}{3} g A^\nu A^\alpha A^\beta \right). \quad (77)$$

In the ‘temporal gauge’ ($A_0 = 0$),

$$\begin{aligned} q &= \frac{g^2}{16\pi^2} \int d^4x \partial^\mu J_\mu = \frac{g^2}{16\pi^2} \Delta_{x_4=\pm\infty} \int d^3x J_0 \\ &= \frac{1}{24\pi^2} \Delta_{x_4=\pm\infty} \int d^3x \epsilon^{ijk} \operatorname{tr} (\partial_i g g^{-1} \partial_j g g^{-1} \partial_k g g^{-1}) = n_+ - n_- . \end{aligned} \quad (78)$$

This means that Euclidean field configurations interpolating between the vacua $|n_+\rangle$, $|n_-\rangle$ at $x_4 = \pm\infty$ have Pontryagin number $q = n_+ - n_-$ and the path integral in Eq.75 should be performed over all these field configurations.

For a given q , there is a lower bound on $S(A)$,

$$S(A) \geq \frac{8\pi^2}{g^2} |q| , \quad (79)$$

which is saturated if and only if $F_{\mu\nu} = \pm \tilde{F}_{\mu\nu}$, i.e, if the configuration is self-dual or self-antidual. For $q=1$, the self-dual classical solution is called instanton⁴⁷ and is given by (in the ‘singular’ gauge)

$$A_{a\mu}(x) = \frac{2\rho^2}{g(x-z)^2} \frac{\eta_{a\mu\nu}(x-z)^\nu}{(x-z)^2 + \rho^2} , \quad (80)$$

where $\eta_{a\mu\nu}$ ($a=1,2,3$; $\mu, \nu=1,2,3,4$) are the t’ Hooft symbols with $\eta_{aij} = \epsilon_{aij}$ ($i, j=1,2,3$), $\eta_{a4i} = -\delta_{ai}$, $\eta_{ai4} = \delta_{ai}$ and $\eta_{a44} = 0$. The instanton depends on four Euclidean coordinates z_μ (its position) and its scale (or size) ρ . Two successive vacua $|n\rangle$, $|n+1\rangle$ are separated by a potential barrier of height $\propto \rho^{-1}$. The Euclidean action of the interpolating instanton is always equal to $8\pi^2/g^2$, but the height of the barrier can be made arbitrarily small since the size ρ of the instanton can be taken arbitrarily large.

We now reintroduce the fermions into the theory and observe⁴⁵ that the B - and L - number currents carry anomalies, i.e.,

$$\partial_\mu J_B^\mu = \partial_\mu J_L^\mu = -n_g \frac{g^2}{16\pi^2} \text{tr}(F_{\mu\nu} \tilde{F}^{\mu\nu}) , \quad (81)$$

where n_g is the number of generations. It is then obvious that the tunneling from $|n_- \rangle$ to $|n_+ \rangle$ is accompanied by a change of the B - and L - numbers, $\Delta B = \Delta L = -n_g q = -n_g(n_+ - n_-)$. Note that i) $\Delta(B - L) = 0$, and ii) for $q=1$, $\Delta B = \Delta L = -3$ which means that we have the annihilation of one lepton per family and one quark per family and color (12-point function).

We, finally, reintroduce the Weinberg-Salam higgs doublet ϕ with its vev given by

$$\langle \phi \rangle = \frac{v}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \end{pmatrix} , \quad v \approx 246 \text{ GeV} . \quad (82)$$

It is then easy to see that the instanton ceases to exist as an exact solution. It is replaced by the so called ‘restricted instanton’⁴⁸ which is an approximate solution for $\rho \ll v^{-1}$. For $|x - z| \ll \rho$, the gauge field configuration of the ‘restricted instanton’ essentially coincides with that of the instanton and the higgs field is

$$\phi(x) \approx \frac{v}{\sqrt{2}} \left(\frac{(x - z)^2}{(x - z)^2 + \rho^2} \right)^{1/2} \begin{pmatrix} 0 \\ 1 \end{pmatrix} . \quad (83)$$

For $|x - z| \gg \rho$, the gauge and higgs fields decay to a pure gauge and the vev in Eq.82 respectively. The action of the ‘restricted instanton’ is $S_{ri} = (8\pi^2/g^2) + \pi^2 v^2 \rho^2 + \dots$, which implies that the contribution of big size ‘restricted instantons’ to the path integral in Eq.75 is suppressed. This justifies *a posteriori* the fact that we restricted ourselves to approximate instanton solutions with $\rho \ll v^{-1}$.

The height of the potential barrier between the vacua $|n\rangle$, $|n+1\rangle$ cannot be now arbitrarily small. This can be understood by observing that the static energy of the ‘restricted instanton’ at $x_4 = z_4$ (λ is the higgs self-coupling),

$$E_b(\rho) \approx \frac{3\pi^2}{g^2} \frac{1}{\rho} + \frac{3}{8} \pi^2 v^2 \rho^2 + \frac{\lambda}{4} \pi^2 v^4 \rho^3 , \quad (84)$$

is minimized for

$$\rho_{\min} = \frac{\sqrt{2}}{gv} \left(\frac{\lambda}{g^2} \right)^{-1/2} \left(\left(\frac{1}{64} + \frac{\lambda}{g^2} \right)^{1/2} - \frac{1}{8} \right)^{1/2} \sim M_W^{-1} , \quad (85)$$

and, thus, the minimal height of the potential barrier turns out to be $E_{\min} \sim M_W/\alpha_W$ (M_W is the weak scale and $\alpha_W = g^2/4\pi$). The static solution which corresponds to the top (saddle point) of this potential barrier is called sphaleron⁴⁹ and is given by

$$\bar{A} = v \frac{f(\xi)}{\xi} \hat{r} \times \bar{\tau} , \quad \phi = \frac{v}{\sqrt{2}} h(\xi) \hat{r} \cdot \bar{\tau} \begin{pmatrix} 0 \\ 1 \end{pmatrix} , \quad (86)$$

where $\xi = 2M_W r$, $\hat{r}$ is the radial unit vector and the 3-vector $\bar{\tau}$ consists of the Pauli matrices. The functions $f(\xi)$, $h(\xi)$, which can be determined numerically, tend to zero as $\xi \rightarrow 0$ and to 1 as $\xi \rightarrow \infty$. The mass of the sphaleron is estimated to be

$$E_{\text{sph}} = \frac{2M_W}{\alpha_W} k, \quad 1.5 \leq k \leq 2.7, \quad \text{for } 0 \leq \lambda \leq \infty , \quad (87)$$

and lies between 10 and 15 TeV.

At zero temperature the tunneling from $|n\rangle$ to $|n+1\rangle$ is utterly suppressed⁴⁵ by the factor $\exp(-8\pi^2/g^2)$. At high temperatures, however, thermal fluctuations over the potential barrier are frequent and this transition can occur⁴¹ with an appreciable rate. For $M_W \lesssim T \lesssim T_c$ (T_c is the critical temperature of the electroweak transition), this rate can be calculated⁴¹ by expanding around the sphaleron (saddle point) solution and turns out to be

$$\Gamma \approx 10^4 n_g \frac{v(T)^9}{T^8} \exp(-E_{\text{sph}}(T)/T) . \quad (88)$$

Assuming that the electroweak phase transition is a second order one, $v(T)$ and $E_{\text{sph}}(T) \propto (1 - T^2/T_c^2)^{1/2}$. One can then show that $\Gamma \gg H$ for temperatures T between ~ 200 GeV and $\sim T_c$. Furthermore, for temperatures above T_c , where the sphaleron solution ceases to exist, it was argued⁴¹ that we still have $\Gamma \gg H$. The overall conclusion is that non-perturbative B - and L - number violating processes are in equilibrium in the universe for $T \gtrsim 200$ GeV. Remember that $B - L$ is conserved by these processes.

Given a primordial L - number density, one can calculate^{42,43} the resulting n_B/s (n_B is the B - number density). In MSSM, the $SU(2)_L$ - instantons produce the effective operator (in symbolic form)

$$O_2 = (qqql)^{n_g} (\tilde{h}^{(1)} \tilde{h}^{(2)}) \tilde{W}^4 , \quad (89)$$

and the $SU(3)_c$ - instantons the operator

$$O_3 = (qqu^c d^c)^{n_g} \tilde{g}^6 , \quad (90)$$

where q, l are the quark, lepton $SU(2)_L$ - doublets respectively, u^c, d^c the up, down type antiquark $SU(2)_L$ - singlets respectively, g, W the gluons and W -bosons and tilde represents their superpartners. We will assume that these interactions together with the usual MSSM interactions are in equilibrium at high temperatures. The equilibrium number density of ultrarelativistic particles $\Delta n \equiv n_{\text{part}} - n_{\text{antipart}}$ is given by

$$\frac{\Delta n}{s} = \frac{15g}{4\pi^2 g_*} \left(\frac{\mu}{T} \right) \epsilon, \quad (91)$$

where g is the number of internal degrees of freedom of the particle under consideration, μ its chemical potential and $\epsilon = 2$ or 1 for bosons or fermions respectively. Each interaction in equilibrium implies that the algebraic sum of the chemical potentials of the particles involved is zero. Solving these constraints, we end up with only two independent chemical potentials, μ_q and $\mu_{\tilde{g}}$, and the baryon and lepton asymmetries can be expressed⁴² in terms of them as follows

$$\begin{aligned} \frac{n_B}{s} &= \frac{30}{4\pi^2 g_* T} (6n_g \mu_q - (4n_g - 9)\mu_{\tilde{g}}), \\ \frac{n_L}{s} &= -\frac{45}{4\pi^2 g_* T} \left(\frac{n_g(14n_g + 9)}{1 + 2n_g} \mu_q + \Omega(n_g) \mu_{\tilde{g}} \right), \end{aligned} \quad (92)$$

where $\Omega(n_g)$ is a known⁴² function. Now soft supersymmetry breaking couplings come in equilibrium at $T \lesssim 10^7$ GeV since their rate $\Gamma_S \approx m_S^2/T \gtrsim H \approx 30 T^2/M_P$ (m_S is the soft supersymmetry breaking scale). In particular, the non-vanishing gaugino mass implies $\mu_{\tilde{g}} = 0$ and Eqs.92 give⁴²

$$\frac{n_B}{s} = \frac{4(1 + 2n_g)}{22n_g + 13} \frac{n_{B-L}}{s}. \quad (93)$$

Equating n_{B-L}/s with the primordial n_L/s , we have $n_B/s = (-28/79)(n_L/s)$, for $n_g = 3$.

12 Supersymmetric Hybrid Inflation and Neutrinos

The supersymmetric model based on the left-right symmetric gauge group G_{LR} , which we discussed in Sec.10, provides a framework within which hybrid inflation, baryogenesis and neutrino oscillations are⁵⁰ closely linked. This scheme, supplemented by a familiar ansatz for the neutrino Dirac masses and mixing of the two heaviest families and with the MSW resolution of the solar neutrino puzzle, implies⁵⁰ relatively stringent restrictions on the mass of the tau-neutrino, m_{ν_τ} , and the $\mu - \tau$ mixing angle, $\theta_{\mu\tau}$.

In order to obtain information about the light neutrino masses and mixing of the two heaviest families, we ignore the first family assuming it has small mixings. The relevant ‘asymptotic’ (at the GUT scale M_X) 2×2 mass matrices are M^l , the mass matrix of charged leptons (l^c, l), M^D , the Dirac mass matrix of neutrinos (ν^c, ν), and M^R , the Majorana mass matrix of ν^c ’s. We shall first diagonalize M^l, M^D :

$$M^l \rightarrow M^{l'} = \tilde{U}^{l^c} M^l U^l = \begin{pmatrix} m_\mu & \\ & m_\tau \end{pmatrix}, \quad (94)$$

$$M^D \rightarrow M^{D'} = \tilde{U}^{\nu^c} M^D U^\nu = \begin{pmatrix} m_2^D & \\ & m_3^D \end{pmatrix}, \quad (95)$$

where the diagonal entries are positive. This gives rise to the ‘Dirac’ mixing matrix $U^{\nu \dagger} U^l$ in the leptonic charged currents. Using the remaining phase freedom, we can bring this matrix to the form

$$U^{\nu \dagger} U^l \rightarrow \begin{pmatrix} \cos \theta^D & \sin \theta^D \\ -\sin \theta^D & \cos \theta^D \end{pmatrix}, \quad (96)$$

where θ^D ($0 \leq \theta^D \leq \pi/2$) is the ‘Dirac’ (not the physical) mixing angle in the 2 – 3 leptonic sector. In this basis, the Majorana mass matrix can be written as $M^R = U^{-1} M_0 \tilde{U}^{-1}$, where $M_0 = \text{diag}(M_2, M_3)$, with M_2, M_3 (both positive) being the two Majorana masses, and U is a unitary matrix which can be parametrized as

$$U = \begin{pmatrix} \cos \theta & \sin \theta e^{-i\delta} \\ -\sin \theta e^{i\delta} & \cos \theta \end{pmatrix} \begin{pmatrix} e^{i\alpha_2} & \\ & e^{i\alpha_3} \end{pmatrix}, \quad (97)$$

with $0 \leq \theta \leq \pi/2$ and $0 \leq \delta < \pi$. The light neutrino mass matrix is

$$m = -\tilde{M}^{D'} \frac{1}{M^R} M^{D'} = \begin{pmatrix} e^{i\alpha_2} & \\ & e^{i\alpha_3} \end{pmatrix} \Psi(\theta, \delta) \begin{pmatrix} e^{i\alpha_2} & \\ & e^{i\alpha_3} \end{pmatrix}, \quad (98)$$

where $\Psi(\theta, \delta)$ depends also on M_2, M_3, m_2^D, m_3^D .

We will denote the two positive eigenvalues of the light neutrino mass matrix by m_2 (or m_{ν_μ}), m_3 (or m_{ν_τ}). Recall that all the quantities here (masses, mixings) are ‘asymptotic’. The determinant and the trace invariance of $m^\dagger m$ provide us with two constraints on the (asymptotic) parameters:

$$m_2 m_3 = \frac{(m_2^D m_3^D)^2}{M_2 M_3}, \quad (99)$$

$$m_2^2 + m_3^2 = \frac{(m_2^{D^2} c^2 + m_3^{D^2} s^2)^2}{M_2^2} + \frac{(m_3^{D^2} c^2 + m_2^{D^2} s^2)^2}{M_3^2} + \frac{2(m_3^{D^2} - m_2^{D^2})^2 c^2 s^2 \cos 2\delta}{M_2 M_3}, \quad (100)$$

where θ, δ are defined in Eq.97, $c = \cos \theta$, $s = \sin \theta$. Note that the phases α_2, α_3 in Eq.98 cancel out in these constraints and, thus, remain undetermined.

The mass matrix m is diagonalized by a unitary rotation V on ν 's:

$$V = \begin{pmatrix} e^{i\beta_2} & \\ & e^{i\beta_3} \end{pmatrix} \begin{pmatrix} \cos \varphi & \sin \varphi e^{-i\epsilon} \\ -\sin \varphi e^{i\epsilon} & \cos \varphi \end{pmatrix}, \quad (101)$$

where $0 \leq \varphi \leq \pi/2$, $0 \leq \epsilon < \pi$. The ‘Dirac’ mixing matrix in Eq.96 is now multiplied by $V^\dagger$ on the left and, after phase absorptions, takes the form

$$\begin{pmatrix} \cos \theta_{23} & \sin \theta_{23} e^{-i\delta_{23}} \\ -\sin \theta_{23} e^{i\delta_{23}} & \cos \theta_{23} \end{pmatrix}, \quad (102)$$

where $0 \leq \theta_{23} \leq \pi/2$, $0 \leq \delta_{23} < \pi$. Here, θ_{23} (or $\theta_{\mu\tau}$) is the physical mixing angle in the 2 – 3 leptonic sector and can be determined from the requirement that its cosine is equal to the modulus of the complex number $\cos \varphi \cos \theta^D + \sin \varphi \sin \theta^D e^{i(\xi - \epsilon)}$, where $-\pi \leq \xi - \epsilon = \beta_2 - \beta_3 - \epsilon \leq \pi$. The phases β_2, β_3 and ξ remain undetermined due to the arbitrariness of α_2, α_3 . Thus, the precise value of θ_{23} cannot be found. However, we can determine the range in which θ_{23} lies: $|\varphi - \theta^D| \leq \theta_{23} \leq \varphi + \theta^D$, for $\varphi + \theta^D \leq \pi/2$.

We now need the asymptotic values of $m_{2,3}^D, \theta^D$. Approximate $SU(4)_c$ -invariance in the up quark and neutrino sectors gives $m_2^D = m_c, m_3^D = m_t, \sin \theta^D = |V_{cb}|$ ‘asymptotically’. Renormalization of light neutrino masses and mixing between M_X and M_Z is also included^{51,52} assuming the MSSM spectrum and large $\tan \beta \approx m_t/m_b$. In the framework of ‘hierarchical’ light neutrino masses ($m_3 \gg m_2 \gg m_1$), the small angle MSW resolution of the solar neutrino puzzle implies⁵³ $1.7 \times 10^{-3} \text{ eV} \lesssim m_2 \lesssim 3.5 \times 10^{-3} \text{ eV}$. Finally, m_3 is restricted by the cosmological bound $m_3 \lesssim 23 \text{ eV}$ (for $h \approx 0.5$).

We are now ready to derive^{34,50} useful restrictions on $M_{2,3}$. Assume that both $M_{2,3} \leq m_{infl}/2$. Then the inflaton predominantly decays to the heaviest of the two. The determinant condition implies that the lowest possible value of the heaviest $M_{2,3}$ is about 10^{11} GeV giving $T_r \gtrsim 10^{10} \text{ GeV}$, in conflict with the gravitino constraint³⁷ (see Sec.10). So we are obliged to require that $1.72 \times 10^{13} \text{ GeV} \approx m_{infl}/2 \leq M_3 \lesssim 2.5 \times 10^{13} \text{ GeV}$, where the upper bound comes from the requirement that the coupling constant of the non-renormalizable term responsible for the mass of the heaviest ν^c does not exceed




Figure 1: The allowed regions in the m_{ν_μ} , m_{ν_τ} plane.

unity. This requirement also implies that $x_Q \lesssim 2.6$, which makes our choice in Sec.9.2, $x_Q \approx 2$, central. In summary, we see that i) M_3 is constraint in a narrow range, and ii) the inflaton decays to the second heaviest right handed neutrino, ν^c , with mass M_2 .

Baryons can be produced, in the present scheme, only via a primordial leptogenesis from the decay of ν^c 's emerging as decay products of the inflaton (see Sec.11.1). The lepton asymmetry is then partially converted into the observed baryon asymmetry of the universe by 'sphaleron' effects as we explained in Sec.11.2. From Eq.71, the lepton asymmetry is

$$\frac{n_L}{s} = \frac{9 T_r}{8\pi m_{infl}} \frac{M_2}{M_3} \frac{c^2 s^2 \sin 2\delta (m_3^{D/2} - m_2^{D/2})^2}{|\langle h^{(1)} \rangle|^2 (m_3^{D/2} s^2 + m_2^{D/2} c^2)} . \quad (103)$$

Renormalization effects should also be included in this formula. Assuming the MSSM spectrum between 1 TeV and M_X , we saw in Sec.11.2 that the observed baryon asymmetry n_B/s is related to n_L/s by $n_B/s = (-28/79)(n_L/s)$.

We will now extract⁵⁰ restrictions on light neutrino masses and mixing. Take a specific value of M_3 (in practice, we take its two extreme values $m_{infl}/2$ or 2.5×10^{13} GeV). For any pair m_2, m_3 , we use the determinant condition




Figure 2: The allowed regions in the ν_μ - ν_τ oscillation plot.

to evaluate M_2 and, thus, T_r . The gravitino constraint³⁷ ($T_r \lesssim 10^9$ GeV) then gives a lower bound in the m_2, m_3 plane. This bound together with the small angle MSW restriction⁵³ on m_2 yields a lower bound on m_3 , namely $m_3 \gtrsim 0.9$ eV (for $M_3 = 2.5 \times 10^{13}$ GeV) or 1.3 eV (for $M_3 = m_{infl}/2$). The trace condition is solved with respect to $\delta = \delta(\theta)$, $0 \leq \theta \leq \pi/2$, which is then substituted in Eq.103 to yield $n_L/s = (n_L/s)(\theta)$. Imposing the ‘low’ deuterium bound⁵⁴ on n_B/s ($0.02 \lesssim \Omega_B h^2 \lesssim 0.03$), we find the range of θ where this bound is satisfied. If such a range exists, we keep m_2, m_3 as satisfying the baryogenesis constraint. This gives an upper bound in the m_2, m_3 plane which together with the MSW restriction on m_2 yields an upper bound on m_3 , namely $m_3 \lesssim 5.1$ eV (for $M_3 = 2.5 \times 10^{13}$ GeV) or 8.8 eV (for $M_3 = m_{infl}/2$). The allowed area in the m_2, m_3 plane is depicted⁵⁰ in Fig.1, where the thick solid (dashed) line corresponds to $M_3 = m_{infl}/2$ ($M_3 = 2.5 \times 10^{13}$ GeV). The overall allowed range for m_{ν_τ} is⁵⁰ $1 \text{ eV} \lesssim m_{\nu_\tau} \lesssim 9 \text{ eV}$, which is interesting since the value of m_{ν_τ} required by the cold plus hot dark matter scenario⁵⁵ for large scale structure formation in the universe is centrally located in this range.

The mixing angle $\theta_{\mu\tau}$ can also be restricted. For every allowed m_2, m_3 pair

and every θ satisfying the baryogenesis constraint, we construct V in Eq.101 and, consequently, φ , ϵ and the allowed range of $\theta_{\mu\tau}$, $|\varphi - \theta^D| \leq \theta_{\mu\tau} \leq \varphi + \theta^D$. The union of all these ranges for all allowed θ 's and m_2 's for a given m_3 gives the range of $\theta_{\mu\tau}$ which is allowed for this value of m_3 . All these ranges for all allowed m_3 's constitute the allowed area on the oscillation diagram, which is depicted⁵⁰ in Fig.2 (notation as in Fig.1) in confrontation to past, ongoing and planned experiments. The central part of this allowed area will be tested by the ongoing Short Baseline Experiments (SBLE) at CERN, NOMAD/CHORUS. Possibly negative result from NOMAD/CHORUS will exclude a significant part of the allowed domains in Figs.1, 2 reducing the upper bound on the tau-neutrino mass, m_{ν_τ} , to 3.7 eV. The new CERN SBLE(TOSCA) together with the new CERN Medium Baseline Experiment (MBLE) ICARUS-JURA (600t,DIS) will cover all our predicted area on the oscillation diagram.

In summary, hybrid inflation, baryogenesis and neutrino oscillations have been linked in the context of a supersymmetric model based on a left-right symmetric gauge group. Our scheme leads to stringent restriction on m_{ν_τ} and $\theta_{\mu\tau}$ to be tested by ongoing and planned experiments. These restrictions are derived by mainly 'physical' arguments (gravitino and baryogenesis constraints) supplemented by a 'minimal' input from fermion mass matrix ansatz (only 3 input parameters) and experiments (MSW resolution of the solar neutrino problem). The choice of the gauge group is crucial since, in this case, ϕ has the quantum numbers of ν^c and, thus, decays to ν^c 's producing an initial lepton asymmetry. As a consequence, the gravitino and baryogenesis constraints restrict the neutrino parameters.

Acknowledgment

This work was supported in part by the European Commission under the Human Capital and Mobility programme, contract number CHRX-CT94-0423.

References

1. For a textbook treatment see e.g., S.Weinberg, *Gravitation and Cosmology* (Wiley, NY, 1972); E.W.Kolb and M.S.Turner, *The Early Universe* (Addison-Wesley, Redwood City, CA, 1990).
2. G.F.Smoot *et al.*, *Astrophys. J. Lett.* **396**, L1 (1992); C.L.Bennett *et al.*, *Astrophys. J. Lett.* **464**, 1 (1996).
3. H.Georgi and S.Glashow, *Phys. Rev. Lett.* **32**, 438 (1974); J.G.Pati and A.Salam, *Phys. Rev. Lett.* **31**, 661 (1973).
4. D.A.Kirzhnits and A.D.Linde, *Phys. Lett.* **42B**, 471 (1972).

5. G.'t Hooft, *Nucl. Phys. B* **79**, 276 (1974); A.Polyakov, *JETP Lett.* **20**, 194 (1974).
6. V.L.Ginzburg, *Sov. Phys. -Solid State* **2**, 1824 (1960).
7. T.W.B.Kibble, *J. Phys. A* **9**, 1387 (1976); *Phys. Reports* **67**, 183 (1980).
8. J.P.Preskill, *Phys. Rev. Lett.* **43**, 1365 (1979).
9. For a pedagogical discussion see e.g., P.J.E.Peebles, *The Large-scale Structure of the Universe* (Princeton Univ. Press, Princeton, 1980); G.Efstathiou, in *The Physics of the Early Universe*, eds. J.A.Peacock, A.F.Heavens and A.T.Davies (Adam-Higler,Bristol, 1990).
10. E.R.Harrison, *Phys. Rev. D* **1**, 2726 (1970); Ya.B.Zeldovich, *Mon. Not. Roy. Astr. Soc.* **160**, 1 (1972).
11. A.H.Guth, *Phys. Rev. D* **23**, 347 (1981).
12. For a textbook introduction see e.g., A.D.Linde, *Particle Physics and Inflationary Cosmology* (Harwood Academic Publishers, London, 1990).
13. A.D.Linde, *Phys. Lett.* **108B**, 389 (1982); A.Albrecht and P.Steinhardt, *Phys. Rev. Lett.* **48**, 1220 (1982).
14. A.D.Linde, *JETP Lett.* **38**, 149 (1983); *Phys. Lett.* **129B**, 177 (1983).
15. R.J.Scherrer and M.S.Turner, *Phys. Rev. D* **31**, 681 (1985).
16. A.Albrecht, P.J.Steinhardt, M.S.Turner and F.Wilczek, *Phys. Rev. Lett.* **48**, 1437 (1982); L.Abbott, E.Farhi and M.Wise, *Phys. Lett.* **117B**, 29 (1982); A.Dolgov and A.D.Linde, *Phys. Lett.* **116B**, 329 (1982).
17. P.J.Steinhardt and M.S.Turner, *Phys. Rev. D* **29**, 2162 (1984).
18. M.S.Turner, *Phys. Rev. D* **28**, 1243 (1983).
19. S.W.Hawking, *Nature* **248**, 30 (1974); *Commun. Math. Phys.* **43**, 199 (1975).
20. G.W.Gibbons and S.W.Hawking, *Phys. Rev. D* **15**, 2752 (1977).
21. T.Bunch and P.C.W.Davies, *Proc. Roy. Soc. London A* **360**, 117 (1978).
22. A.Vilenkin and L.Ford, *Phys. Rev. D* **26**, 1231 (1982); A.D.Linde, *Phys. Lett.* **116B**, 335 (1982); A.A.Starobinsky, *Phys. Lett.* **117B**, 175 (1982).
23. W.Fischler, B.Ratra and L.Susskind, *Nucl. Phys. B* **259**, 730 (1985).
24. J.M.Bardeen, P.J.Steinhardt and M.S.Turner, *Phys. Rev. D* **28**, 679 (1983); for a review see e.g., V.F.Mukhanov, H.A.Feldman and R.H.Brandenberger, *Phys. Reports* **215**, 203 (1992).
25. G.Lazarides and Q.Shafi, *Phys. Lett. B* **308**, 17 (1993).
26. J.M.Bardeen, *Phys. Rev. D* **22**, 1882 (1980); H.Kodama and M.Sasaki, *Prog. Theor. Phys.* **72**, 1 (1984).
27. L.F.Abbott and R.K.Schaefer, *Astrophys. Journal* **308**, 546 (1986); R.K.Schaefer, *Phys. Rev. D* **6**, 2075 (1987).
28. R.K.Sachs and A.M.Wolfe, *Astrophys. Journal* **147**, 73 (1967).
29. V.A.Rubakov, M.Sazhin and A.Veryaskin, *Phys. Lett. B* **115**, 189 (1982);

- R.Fabbri and M.Pollock, *Phys. Lett.* **125B**, 445 (1983); B.Allen, *Phys. Rev. D* **37**, 2078 (1988); L.Abbott and M.Wise, *Nucl. Phys. B* **244**, 541 (1984); M.White, *Phys. Rev. D* **46**, 4198 (1992).
30. A.D.Linde, *Phys. Lett. B* **259**, 38 (1991); *Phys. Rev. D* **49**, 748 (1994).
 31. E.J. Copeland, A.R. Liddle, D.H. Lyth, E.D. Stewart and D. Wands, *Phys. Rev. D* **49**, 6410 (1994).
 32. G.Lazarides and C.Panagiotakopoulos, *Phys. Rev. D* **52**, R559 (1995).
 33. G.Dvali, Q.Shafi and R.K.Schaefer, *Phys. Rev. Lett.* **73**, 1886 (1994).
 34. G.Lazarides, R.K.Schaefer and Q.Shafi, *Phys. Rev. D* **56**, 1324 (1997).
 35. G.Lazarides and N.Tetradis, *Phys. Rev. D* **58**, 123502 (1998); E.D. Stewart, *Phys. Rev. D* **51**, 6847 (1995).
 36. G.Dvali, G.Lazarides and Q.Shafi, *Phys. Lett. B* **424**, 259 (1998).
 37. M.Yu.Khlopov and A.D.Linde, *Phys. Lett.* **138B**, 265 (1984); J.Ellis, J.E.Kim and D.Nanopoulos, *Phys. Lett.* **145B**, 181 (1984); M.Kawasaki and T.Moroi, *Prog. Theor. Phys.* **93**, 879 (1995).
 38. Q.Shafi and X.M.Wang, unpublished(1990); L.Ibanez and G.G.Ross, *Nucl. Phys. B* **368**, 3 (1992); G.Lazarides, C.Panagiotakopoulos and Q.Shafi, *Phys. Lett. B* **315**, 325 (1993), (E) *ibid.* **317**, 661 (1993).
 39. G.Lazarides, C.Panagiotakopoulos and Q.Shafi, *Phys. Rev. Lett.* **58**, 1707 (1987).
 40. M.Fukugita and T.Yanagida, *Phys. Lett. B* **174**, 45 (1986); G.Lazarides and Q.Shafi, *Phys. Lett. B* **258**, 305 (1991).
 41. S.Dimopoulos and L.Susskind, *Phys. Rev. D* **18**, 4500 (1978); V.Kuzmin, V.Rubakov and M.Shaposhnikov, *Phys. Lett. B* **155**, 36 (1985); P.Arnold and L.McLerran, *Phys. Rev. D* **36**, 581 (1987).
 42. L.Ibáñez and F.Quevedo, *Phys. Lett. B* **283**, 261 (1992).
 43. J.A.Harvey and M.S.Turner, *Phys. Rev. D* **42**, 3344 (1990).
 44. L.Covi, E.Roulet and F.Vissani, *Phys. Lett. B* **384**, 169 (1996); W.Buchmüller and M.Plümacher, *Phys. Lett. B* **431**, 354 (1998).
 45. G.'t Hooft, *Phys. Rev. Lett.* **37**, 8 (1976); *Phys. Rev. D* **14**, 3432 (1976).
 46. C.Callan, R.Dashen and D.Gross, *Phys. Lett.* **63B**, 334 (1976); R.Jackiw and C.Rebbi, *Phys. Rev. Lett.* **37**, 172 (1976).
 47. A.A.Belavin, A.Polyakov, A.Schwartz and Y.Tyupkin, *Phys. Lett.* **59B**, 85 (1975).
 48. I.Affleck, *Nucl. Phys. B* **191**, 429 (1981).
 49. N.S.Manton, *Phys. Rev. D* **28**, 2019 (1983); F.R.Klinkhamer and N.S.Manton, *Phys. Rev. D* **30**, 2212 (1984).
 50. G.Lazarides, Q.Shafi and D.N.Vlachos, *Phys. Lett. B* **427**, 53 (1998).
 51. K.S.Babu, C.N.Leung and J.Pantaleone, *Phys. Lett. B* **319**, 191 (1993).

- 52. K.S.Babu and Q.Shafi, *Phys. Rev. D* **47**, 5004 (1993); *Phys. Lett. B* **357**, 365 (1995).
- 53. See e.g., A.Smirnov, *hep-ph/9611465* and references therein.
- 54. D.Tytler, X.M.Fan and S.Burles, *Nature* **381**, 207 (1996); S.Burles and D.Tytler, *Astrophys. Journal* **460**, 584 (1996).
- 55. Q.Shafi and F.W.Stecker, *Phys. Rev. Lett.* **53**, 1292 (1984). For a recent review and other references see Q.Shafi and R.K.Schaefer, *hep-ph/9612478*.